

## The Physicochemical Properties of Starch from *Tongil*-type Rice Varieties

Jong-Hee Shin<sup>1,†</sup>, Chae-Min Han<sup>1</sup>, Young-Un Song<sup>1</sup>, Sang-Kuk Kim<sup>2</sup>, and Jung-Gi Ryu<sup>2</sup>

**ABSTRACT** Starch characteristics and pasting properties of *Tongil*-type rice varieties with different amylose content were analyzed. Three different *Tongil*-type rice varieties and one *Japonica*-type rice variety were examined for their properties. *Tongil*-type rice varieties have longer panicles and higher rice yield (721-765 kg per 10a) than Boramchan, a *Japonica*-type high-yield rice variety. The protein content of the *Tongil*-type rice variety was higher than that of Boramchan. *Japonica*-type rice varieties had lower amylose content than *Japonica*-type rice varieties, except for Amimyeon (Milyang355). Amimyeon had higher protein content than the other varieties, and its amylose content was particularly high at 39.2%. The distribution of starch granule sizes of the four varieties was similar, and the particle size corresponding to D50 was approximately 87.8-81.9  $\mu\text{m}$ . The pasting properties of rice flour varied among varieties. The Dasanbyeon and Hanarum2 amylogram patterns were similar. These two varieties had a higher peak viscosity (PV) and lower setback (SB) than Boramchan. In the case of Amimyeon, the hot paste viscosity (HPV), cool paste viscosity (CPV), and SB were significantly higher than those of the other *Toingil*-type varieties, and the breakdown (BD) was very low, showing pasting properties significantly different from that of the other varieties. The onset ( $T_o$ ), peak ( $T_p$ ), and conclusion temperature ( $T_c$ ) of gelatinization of rice flour from Amimyeon were lower than those of the other tested varieties. In addition, the gelatinization temperature, measured using differential scanning calorimetry (DSC), and BD, measured using a rapid visco analyzer (RVA) were low in Amimyeon. Amylose content showed a high positive correlation with pasting time (PTi), HPV, and SB and a negative correlation with PV and BD.

**Keywords** : gelatinization, pasting property, rice starch, *Tongil*-type

The rice protein, amylose, Mg, and K content is known as chemical factor affecting the taste of rice. The content of amylose and protein is widely used as a criterion for determining the rice taste. The lower the amylose and protein content, the better the rice tastes (Kim *et al.*, 1991). However, among rice varieties with similar amylose and protein content, there are varieties that show differences in rice taste, especially *Tongil*-type rice varieties. Although the *Tongil*-type rice varieties have similar or lower amylose content than general *Japonica*-type rice varieties, the panel's taste valuation results are not good (Lee *et al.*, 2014). Since there are limitations in explaining the taste of rice only with amylose or protein content, various attempts have been to interpret the factors related to rice taste, such as starch studies using electron microscopy (Kum *et al.*,

2004; Kim *et al.*, 2005), fat content of rice endosperm (Yoon *et al.*, 2012), and storage protein characteristics (Kwak *et al.*, 2016). As a result of previous studies, the proteins surrounding starch granules is directly involved in water permeability, starch gelatinization, and suppression of swelling during the cooking process, and affects the elasticity and viscosity of rice (Son *et al.*, 2002; Choi, 2002). In addition,  $\alpha$ -1, which are subunits of acidic glutelin, albumin, and globulin affects the rice taste of *Tongil*-type rice varieties (Kwak *et al.*, 2016).

Starch is the main source of energy stored in cereal grains. The amount of starch in grains varies, but typically accounts for 60-75% of grain weight and provides 70-80% of calories consumed by humans worldwide (Thomas & Atwell, 1999). Starch is the most abundant component in rice grains, making

<sup>1</sup>Agriculture Researcher, Division of Crop Research, Gyeongsangbuk-do Provincial Agricultural Research and Extension Services, Daegu 41404, Republic of Korea

<sup>2</sup>Agriculture Senior Researcher, Division of Crop Research, Gyeongsangbuk-do Provincial Agricultural Research and Extension Services, Daegu 41404, Republic of Korea

<sup>†</sup>Corresponding author: Jong-Hee Shin; (Phone) +82-53-320-0271; (E-mail) [szzong91@korea.kr](mailto:szzong91@korea.kr)

<Received 7 June, 2023; Revised 12 August, 2023; Accepted 14 August, 2023>

up approximately 90% of the dry weight of milled rice grains (Fitzerald *et al.*, 2004). Starch determines the eating and cooking quality of rice grains, at least contributes to them through interactions with other components in the rice endosperm (proteins, lipids, water) or through interactions with other ingredients used to process the rice (Fitzerald *et al.*, 2004). The content and structure of amylose and amylopectin affect the architecture of the starch granule, gelatinization and pasting profiles, and textural attributes.

The physicochemical properties of rice starches are highly dependent on the rice variety, environment, cultural practices, and extraction conditions. Many studies have revealed that the physicochemical characteristics such as the amylose/amylopectin ratio and granule size are responsible for the properties of starch (Singh *et al.*, 2003). Cereals high in amylose content and resistant starch offer potential health benefits. Resistant starch is kind of starch or starch products that are not digestible and absorbed in the stomach or small intestine and passed on directly to the large intestine (Asp, 1992). A cereal grain higher in amylose content is always a good source of resistant starch (Jiang *et al.*, 2010). Cereals high in resistant starch are benefit to improve human health and to reduce the risk of those serious non-infectious disease (Regina *et al.*, 2006).

In this study, the starch characteristics and pasting properties of *Tongil*-type rice varieties with different amylose content were compared with Boramchan, a *Japonica*-type rice variety, and analyzed.

## MATERIAL AND METHOD

### Sample preparation and experimental design

The varieties used in the experiments includes three *Tongil*-type rice variety (Dasanbyeon, Hanarum2, Amimyeon) and one *Japonica*-type rice variety (Boramchan). In 2020~2022, these varieties were transplanted after puddling a designated experimental plot in May 30<sup>th</sup> at a paddy field of Gyeongsangbuk -do agricultural Research & Extension Services. The planting distance was 30×12 cm, the amount of the fertilizer applied was N-P<sub>2</sub>O<sub>5</sub>-K<sub>2</sub>O=18-9-11 kg 10a<sup>-1</sup>, and the N fertilizer was split applied as a basal, tillering, panicle initiation application with a ratio of 50-25-25. When the grain moisture lowered to 15%, after removing the rice husks with a laboratory milling machine (SY94+RTA2+2400, Ssangyong, Korea), pulverizing them

using a food grinder (HMF-2100S, Hanil Electronics, Korea). Then, flour from all varieties was passed through a 100 mesh sieve and used as test sample.

### Milled rice quality

The husked grain was milled by uniformly placing pressure (SY21-NSART1003, Ssangyong, Korea), achieving 90% milled rice per brown rice ratio. Milled rice quality such as head rice and crushed rice ratio were measured in a Grain Analyzer (Cervitec Grain Inspector 1625, Foss, Sweden). Rice endosperm protein and amylose contents were measured using Near-Infrared Grain Tester (Infratec 1241, Foss, Sweden).

### Starch isolation

Starch was isolated through alkaline treatment after soaking the rice grains in water (Yamamoto & Shirakawa, 1999). After drying the soaked rice, it was dried and pulverized using a blender, and steeped repeatedly in 0.2% NaOH solution. Thereafter, the precipitate was collected, thoroughly washed with deionized water, neutralized with 1N HCl, and centrifuged at room temperature at 1,300 × g for 10 min (VS-21SMT, Vision Scientific Co., Ltd, Korea). The isolated starch was dried at room temperature and sieved with 100 mesh screen.

### Granule size distribution and damaged starch contents

A laser diffraction particle size analyzer (Malvern Mastersizer 2000, Malvern Instruments Ltd., UK) was used to measure the granule size distribution of flour from four different rice varieties. Powder samples were then immersed in ethyl alcohol for 30 s after sonication.

Damaged starch content was measured using the Megazyme kit (Megazyme International Ireland, Wicklow Ireland) according to the American Association of Cereal Chemists (AACC) method 76-31 (AACC, 76-31).

### Gelatinization properties

#### Starch pasting properties

Starch pasting properties of rice flours were measured with a Rapid Visco Analyzer (RVA, Model 4, Newport Scientific, Australia). RVA characterization included initial pasting temperature, peak viscosity, trough viscosity, final viscosity, breakdown (BD), and setback (SB) from the RVA viscogram.

**Table 1.** Growth and yield components of *Tongil*-type rice varieties.

Variety	Heading date (M.D)	Culm length (cm)	Panicle length (cm)	Panicle number (No/plant)	Spikelet number (No/panicle)	Grain ripening (%)	1,000 grain weight of brown rice (g)
Dasanbyeo	8. 6. $\pm$ 2.7 <sup>†</sup>	75 $\pm$ 5.1	23 $\pm$ 1.0	13 $\pm$ 0.5	112 $\pm$ 16.8	80.3 $\pm$ 5.40	25.8 $\pm$ 0.40
Hanarum2	8. 8. $\pm$ 0.0	76 $\pm$ 3.5	23 $\pm$ 1.0	14 $\pm$ 0.9	98 $\pm$ 7.8	83.0 $\pm$ 2.34	24.5 $\pm$ 0.00
Amimyeon	8.13. $\pm$ 2.7	72 $\pm$ 1.5	22 $\pm$ 1.0	14 $\pm$ 1.0	111 $\pm$ 6.5	91.2 $\pm$ 1.70	21.2 $\pm$ 0.12
Boramchan <sup>††</sup>	8.15. $\pm$ 4.6	71 $\pm$ 4.6	21 $\pm$ 0.6	15 $\pm$ 0.5	93 $\pm$ 7.8	83.1 $\pm$ 9.12	22.9 $\pm$ 1.33

<sup>†</sup>All data represent the mean  $\pm$  SD of three measurements.

<sup>††</sup>*Japonica*-type ‘Boramchan’ was tested as a comparative variety.

**Table 2.** Protein, amylose contents, and milled rice quality of *Tongil*-type rice varieties.

Variety	Protein (%)	Amylose (%)	Head rice (%)	Crushed rice (%)
Dasanbyeo	7.5 $\pm$ 0.02 <sup>b†</sup>	17.6 $\pm$ 0.18 <sup>c</sup>	30.1 $\pm$ 8.43 <sup>c</sup>	66.5 $\pm$ 9.08 <sup>a</sup>
Hanarum2	7.6 $\pm$ 0.06 <sup>b</sup>	17.8 $\pm$ 0.16 <sup>c</sup>	55.3 $\pm$ 1.79 <sup>b</sup>	41.3 $\pm$ 1.67 <sup>b</sup>
Amimyeon	8.7 $\pm$ 0.06 <sup>a</sup>	39.2 $\pm$ 0.52 <sup>a</sup>	29.2 $\pm$ 5.08 <sup>c</sup>	62.0 $\pm$ 6.60 <sup>a</sup>
Boramchan <sup>††</sup>	6.8 $\pm$ 0.05 <sup>c</sup>	19.0 $\pm$ 0.04 <sup>b</sup>	81.2 $\pm$ 1.82 <sup>a</sup>	13.6 $\pm$ 1.53 <sup>c</sup>

<sup>†</sup>All data represent the mean  $\pm$  SD of three measurements. Different letters in the same column indicate significant differences among the samples according to Duncan’s multiple range test ( $P < 0.05$ ).

<sup>††</sup>*Japonica*-type ‘Boramchan’ was tested as a comparative variety.

### Differential scanning calorimetry (DSC)

The gelatinization properties of starch were evaluated using Differential scanning calorimetry (DSC). Briefly, 3.0 mg of rice flour and deionized water (1:2, v/v) were left for 1 h and then heated from 30°C to 100°C at a rate of 10°C/min using a DSC (DSC 8500, Perkin Elmer, Waltham, MA, USA). From this peak, the gelatinization onset temperature ( $T_o$ ), gelatinization peak temperature ( $T_p$ ), gelatinization conclusion temperature ( $T_c$ ), and gelatinization enthalpy ( $\Delta H$ ) were measured in triplicate.


### Statistical analysis

One-way ANOVA of the average values was performed to identify significant differences between-groups ( $P < 0.05$ ). Duncan’s multiple range test was also performed to identify differences between treatment using R statistical software (version 3.6.2). All tests were performed in triplicate.

## RESULT AND DISCUSSION

### Growth, yield and milled rice quality

*Tongil*-type rice varieties had longer panicles and higher rice yields (721-765 kg 10a<sup>-1</sup>) than Boramchan, a high-quantity

**Fig. 1.** Comparison of milled rice yield of *Tongil*-type rice varieties.

Bar represent the mean  $\pm$  SD of three measurements. *Japonica*-type ‘Boramchan’ was tested as a comparative variety.

*Japonica*-type rice variety (Table 1, and Fig. 1). The protein content of *Tongil*-type rice varieties was relatively higher than that of Boramchan, a *Japonica*-type rice variety, and the head rice ratio was low. *Tongil*-type rice varieties, except Amimyeon (Milyang355), had lower amylose content than Boramchan (Table 2). Amimyeon had higher protein content than the other varieties, and its amylose content was particularly high at 39.2%.

Amimyeon is a new rice variety developed by Rural Development Administration in 2022. In general, *Tongil*-type rice varieties have higher yield than *Japonica*-type rice varieties and are characterized by long grains (Sim *et al.*, 2018).


**Granule size distribution and damaged starch content**

The size distribution of rice flour granules directly affects the quality of final products, including bread and noodles, due to changes in the gelatinization properties (Hallick & Kelly, 1992) and gel consistency (Cagampang *et al.*, 1973) of rice flours (Evers & Stevens, 1985). The particle size distribution and damaged starch contents of each variety is reported in Table 3. The distribution of starch granule size of four varieties used in the experiment was similar, and the particle size corresponding to D50 was approximately 87.8-81.9 μm. The damaged starch content was 3.2-4.5%. Dasanbyeo had a relatively small amount of 3.2% damaged starch. Damaged starch can be used as a substrate for starch hydrolases, but excessive proportions of damaged starch can affect dough quality negatively.

**Gelatinization properties**

**Pasting properties**

The pasting properties of rice flour varied among varieties (Fig. 2, and Table 4). Pasting is the phenomenon following



**Fig. 2.** Comparison of pasting patterns (RVA profiles) of *Tongil*-type rice varieties. \**Japonica*-type ‘Boramchan’ was tested as a comparative variety.

**Table 3.** Particle size distribution (D10, D50, and D90) of *Tongil*-type rice varieties.

Variety	Particle size (μm)			Damaged starch (%)
	D10 <sup>†</sup>	D50	D90	
Dasanbyeo	21.6 ± 1.33 <sup>ns††</sup>	88.3 ± 2.48 <sup>ns</sup>	180.8 ± 2.88 <sup>ns</sup>	3.2 <sup>b</sup>
Hanarum2	22.7 ± 2.45	91.9 ± 4.83	185.1 ± 6.45	4.5 <sup>a</sup>
Amimyeon	21.6 ± 1.13	88.1 ± 2.51	179.3 ± 3.41	4.0 <sup>a</sup>
Boramchan <sup>††</sup>	21.1 ± 1.74	87.8 ± 2.47	178.2 ± 2.21	3.9 <sup>a</sup>

<sup>†</sup>D10, D50, and D90 represent 10%, 50%, and 90% of the cumulative particle size distribution, respectively.

<sup>††</sup>All data represent the mean ± SD of three measurements. Different letters in the same column of each variety indicate significant differences among the samples according to Duncan’s multiple range test (*P* < 0.05).

<sup>†††</sup>*Japonica*-type ‘Boramchan’ was tested as a comparative variety.

**Table 4.** RVA pasting parameters of *Tongil*-type rice varieties.

Variety	Pasting time (min)	Pasting temp. (°C)	Viscosity (cP)				
			PV <sup>††</sup>	HPV	CPV	BD	SB
Dasanbyeo	4.0 ± 0.08 <sup>a†</sup>	85.1 ± 0.97 <sup>ab</sup>	3,285 ± 91.9 <sup>a</sup>	1,760 ± 47.7 <sup>b</sup>	2,953 ± 46 <sup>b</sup>	1,525 ± 51 <sup>b</sup>	-332 ± 69 <sup>c</sup>
Hanarum2	3.9 ± 0.00 <sup>b</sup>	84.5 ± 0.00 <sup>b</sup>	3,368 ± 23.7 <sup>a</sup>	1,711 ± 93.5 <sup>b</sup>	2,798 ± 118 <sup>b</sup>	1,657 ± 71 <sup>a</sup>	-571 ± 96 <sup>d</sup>
Amimyeon	4.1 ± 0.04 <sup>a</sup>	85.9 ± 0.49 <sup>a</sup>	2,621 ± 35.5 <sup>c</sup>	2,296 ± 24.7 <sup>a</sup>	4,439 ± 99 <sup>a</sup>	325 ± 11 <sup>d</sup>	1,818 ± 115 <sup>a</sup>
Boramchan <sup>†††</sup>	3.5 ± 0.64 <sup>c</sup>	84.2 ± 1.63 <sup>b</sup>	2,961 ± 8.6 <sup>b</sup>	1,760 ± 64.8 <sup>b</sup>	2,849 ± 60 <sup>b</sup>	1,201 ± 56 <sup>c</sup>	-112 ± 52 <sup>b</sup>

<sup>†</sup>All data represent the mean ± SD of three measurements. Different letters in the same column of each variety indicate significant differences among the samples according to Duncan’s multiple range test (*P* < 0.05).

<sup>††</sup>PV, HPV, and CPV are the peak viscosity, hot paste viscosity, and cool paste viscosity, respectively. BD and SB are the breakdown and setback, respectively. BD is equal to the difference between PV and HPV, consistency is equal to the difference between the final viscosity and HPV, and SB is equal to the difference between the CPV, and PV.

<sup>†††</sup>*Japonica*-type ‘Boramchan’ was tested as a comparative variety.

**Table 5.** DSC parameters of *Tongil*-type rice varieties.

Variety	$T_o^\dagger$ (°C)	$T_p$ (°C)	$T_c$ (°C)	$\Delta T$ ( $T_c-T_o$ )	$\Delta H$ (J/g)
Dasanbyeo	$63.2 \pm 0.14^{\dagger\dagger}$	$68.4 \pm 0.42$	$73.8 \pm 0.99$	$10.6 \pm 1.06$	$5.7 \pm 2.94$
Hanarum2	$62.8 \pm 0.14$	$69.8 \pm 0.32$	$75.3 \pm 0.48$	$12.4 \pm 0.34$	$8.7 \pm 0.91$
Amimyeon	$59.7 \pm 0.07$	$66.9 \pm 0.23$	$73.3 \pm 0.56$	$13.6 \pm 0.53$	$7.1 \pm 0.35$
Boramchan <sup>†††</sup>	$63.5 \pm 0.33$	$69.3 \pm 0.02$	$75.0 \pm 0.38$	$11.6 \pm 0.71$	$8.0 \pm 1.39$

$T_o$ ,  $T_p$ , and  $T_c$  are the temperatures of the onset, peak, and conclusion of gelatinization, respectively.  $\Delta T$  ( $T_c-T_o$ ) is the temperature range of gelatinization, and  $\Delta H$  is the enthalpy change of gelatinization.


<sup>††</sup>All data represent the mean  $\pm$  SD of three measurements.

<sup>†††</sup>*Japonica*-type ‘Boramchan’ was tested as a comparative variety.

gelatinization in the dissolution of a starch and involves granular swelling, exudation of molecular components from the granules, and eventually, total disruption of the granules (Thomas & Atwell, 1999). In general, *Japonica*-type varieties with a high taste value are known to have low pasting temperatures, HPV and BD, and low final viscosity (Choi *et al.*, 2006). The Dasanbyeo and Hanarum2 amylogram patterns were similar. These two varieties had a higher PV, and lower SB than Boramchan, a *Japonica*-type rice variety. In the case of Amimyeon, the HPV, CPV, and SB were significantly higher than those of the other *Toingil*-type varieties, and the BD was very low, showing pasting properties significantly different from that of the other varieties. This result was similar to the report of Sim *et al.* (2018) comparing the pasting properties of *Tongil*-type rice and *Japonica*-type rice flour for rice porridge and found that the *Tongil*-type rice varieties had relatively high BD. In the case of Amimyeon with high amylose content, the pasting pattern was markedly different from other rice varieties.

#### Thermal properties

Differential scanning calorimetry (DSC) was used for real-time heat flow analysis to determine how transplanting time affected the thermal properties of three different *Tongil*-type rice varieties (Fig. 3, and Table 5). The  $T_o$ , and  $T_p$  of gelatinization of rice flour of Amimyeon were lower than those of the other tested varieties. In addition, the gelatinization temperature, measured by DSC and BD in RVA were low in Amimyeon. DSC is a thermal analysis that can thermodynamically describe the gelatinization process by measuring enthalpy from an endothermic reaction of gelatinization pasting because it can measure heat absorption or release by chemical reactions during phase changes,



**Fig. 3.** Comparison of DSC curves (Thermal properties) of *Tongil*-type rice varieties.

<sup>†</sup>*Japonica*-type ‘Boramchan’ was tested as a comparative variety.

such as melting (Lee *et al.*, 1993). In the case of Dasanbyeo the lowest gelatinization enthalpy was shown among tested varieties, indicating that lower gelatinization enthalpy required less energy to disrupt (i.e., melt) starch crystallinity.

#### Correlation between amylose content and pasting properties

The amylose content in milled rice had a significant effect on rice starch characteristics (Table 6). The correlation between amylose content and pasting properties of *Tongil*-type rice varieties was analyzed. The amylose content showed a high positive correlation with pasting time (PTi), HPV, and SB and a negative correlation with PV and BD. The amylose content also

**Table 6.** Correlation between rice starch properties and amylose content of three *Tongil*-type rice varieties.

Starch granule size (D50)	RVA pasting parameters						
	PTi <sup>†</sup>	PTe	PV	HPV	CPV	BD	SB
-0.53 <sup>**††</sup>	0.86 <sup>*</sup>	0.90 <sup>**</sup>	-0.99 <sup>**</sup>	1.00 <sup>*</sup>	0.99	-1.00 <sup>*</sup>	0.99

<sup>†</sup>PTi, PTe, PV, HPV, CPV, BD, and SB are the pasting time, pasting temperature, peak viscosity, hot paste viscosity, cool paste viscosity, breakdown, and setback, respectively.

<sup>††</sup>The values are the correlation coefficients between the amylose content and each starch property.

\*, \*\* Significant at 5% and 1% levels compared with amylose content and each starch property.

showed a negative correlation with the starch granule size with a correlation coefficient of 0.53. It is known that the higher the amylose content of rice flour, the lower the PV and the higher the SB (Rho & Ahn, 1989; Yoon *et al.*, 2011).

## CONCLUSION

*Tongil*-type rice varieties had higher protein and lower amylose content than *Japonica*-type rice variety. except Amimyeon (Milyang355). Amimyeon had higher protein content than the other varieties, and the amylose content was particularly high at 39.2%. Dasanbyeon and Hanarum2 have higher PV and lower SB than *Japonica*-type rice variety. These are properties that adversely affect the taste of *Japonica*-type rice variety (Choi *et al.*, 2006). In the case of Amimyeon with high amylose content, the pasting pattern was markedly different from other rice varieties. *Tongil*-type rice varieties that have high yields and high amylose content are recognized as proper processing varieties due to their suitable starch properties.

## REFERENCES

- Asp, N. G. 1992. Resistant starch: proceedings from the 2nd plenary meeting of EURESTA: European FLAIR Concerted Action no. 11(COST 911): physiological implications of the consumption of resistant starch in man. *European Journal of Clinical Nutrition* 46 : S1-S148.
- Cagampan, G. B., C. M. Perez, and B. O. Juliano. 1973. A gel consistency test for eating quality of rice. *Journal of the Science of Food and Agriculture* 24(12) : 1589-1594.
- Choi, B. K., J. S. Kum, H. Y. Lee, and J. D. Park. 2006. Physico-chemical properties of black rice flours affected by milling conditions. *Korean Journal of Food Science Technology* 38 : 751-855.
- Choi, H. 2002. Current status and perspectives in varietal improvement of rice cultivars for high-quality and value-added products. *Korean Journal of Crop Science* 47 : 15-32.
- Evers, A. D. and D. J. Stevens. 1985. Starch damage. *Advances in Cereal Science and Technology* 7 : 321-349.
- Fitzerald, H. A., M. Chern, R. Navarre, and P. C. Ronald. 2004. Over expression of (At) NPRI in rice leads to a BTH-and environment-induced lesion-mimic/cell death phenotype. *Molecular Plant-Microbe Interactions* 17(2) : 140-151.
- Hallick, J. V. and V. J. Kelly. 1992. Gelatinization and pasting characteristics of rice varieties as related to cooking behaviour. *Cereal Chemistry* 36 : 91-98.
- Jiang, H., J. Lio, M. Blanco, M. Campbell, and J. L. Jane. 2010. Resistant-starch formation in high-amylose maize starch during kernel development. *Journal of Agricultural and Food Chemistry* 58 : 8043-8047.
- Kim, K. S., K. J. Kang, and S. K. Kim. 1991. Relationship between hot water solubles of rice and texture of cooked rice. *Korean Journal of Food Science Technology* 23(5) : 498-502.
- Kim, Y. D., U. G. Ha, Y. C. Song, J. H. Cho, E. I. Yang, and J. K. Lee. 2005. Palatability evaluation and physical characteristics of cooked rice. *Korean Journal of Crop Science* 50 : 24-28.
- Kum, K. S., H. J. Kang, I. K. Hwang, H. Hwang, T. Y. Kim, and H. C. Choi. 2004. Comparative ultrastructure of Ilpumbyeon, a high-quality japonica rice, and its mutant, Suwon 464 : Scanning and transmission electron microscopy studies. *Journal of Agricultural and Food Chemistry* 52(12) : 3876-3883.
- Kwak, J., J. S. Lee, M. R. Yoon, M. J. Kim, A. Chun, and C. K. Lee. 2016. Characteristics of seed storage protein affecting the eating quality of *Japonica* and *Tongil*-type rice. *Korean Journal of Crop Science* 61(4) : 227-234.
- Lee, B. Y., C. K. Mok, and C. H. Lee. 1993. Comparison of differential scanning calorimetry with enzymatic method for the determination of gelatinization degree of corn starch. *Korean Journal of Food Science and Technology* 25(4) : 400-403.
- Lee, J. S., J. K. Lee, M. R. Yoon, J. Kwak, A. Chun, K. H. Kang, J. U. Jeung, J. Chun, and B. K. Kim. 2014. Varietal preference of grain appearance and sensory of African on rice ecotypes. *Korean J. Int. Agric.* 26(3) : 232-237.
- Regina, A., A. Bird, D. Topping, S. Bowdem, J. Freeman, T. Barsby, A. Kosar-Hashemi, Z. Li, S. Rahman, and M. Morell. 2006. High-amylose wheat generated by RNA interference improves indices of large-bowel health in rats. *Proceedings of*

- the National Academy of Science. U.S.A. 103 : 3546-3551.
- Rho, E. S. and S. Y. Ahn. 1989. Texture of cooked rice and molecular weight distribution of rice amylose. *Korean Journal of Food Science Technology* 21 : 486-491.
- Sim, E. Y., J. Y. Lee, J. H. Cho, M. R. Yoon, J. Kwak, N. G. Kim, Y. H. Jeon, C. K. Lee, J. S. Lee, and H. C. Hong. 2018. Quality characteristics of porridge made from different Korean rice varieties including high yield tongil-type rice. *Korean Journal of Food Preservation* 25(6) : 651-658.
- Singh, N., J. Singh, L. Kaur, N. S. Sodhi, and B. S. Gill. 2003. Morphological, thermal and rheological properties of starches from different botanical sources. *Food Chemistry* 81(2) : 219-231.
- Son, J. R., J. H. Kim, J. I. Lee, Y. H. Youn, J. K. Kim, H. G. Hwang, and H. P. Moon. 2002. Trend and further research of rice quality evaluation. *Korean Journal of Crop Science* 47 : 33-54.
- Thomas, D. J. and W. A. Atwell. 1999. *Starches*. St. Paul (Minnesota): Eagan Press.
- Yamamoto, A. and K. Shirakawa. 1999. Annealing of long-term stored rice grains improves gelatinization properties. *Cereal Chemistry* 76(5) : 646-649.
- Yoon, M. R., A. R. Chun, S. K. Oh, S. H. Ko, D. J. Kim, H. C. Hong, I. S. Choi, and J. H. Lee. 2011. Physicochemical properties of endosperm starch and breadmaking quality of rice cultivars. *Korean Journal of Crop Science* 56 : 219-225.
- Yoon, M. R., S. C. Lee, and M. Y. Kang. 2012. The lipid composition of rice cultivars with different eating qualities. *Journal of Applied Biological Chemistry* 55 : 291-295.